

STARICHKOV, Vladimir Semenovich; OSTROVA, I.M., red.; MALINOVSKIY,
Yu.F., red.; KYABOV, N.F., nauchnyy red.; PERSON, M.N.,
tekh. red.

[An aid for the master machinist] V pomoshch' masteru-
slesariu; (al'bom). Moskva, Proftekhizdat, 1961. 225 p.
(MIRA 15:8)

(Metal cutting)

DUDKIN, M.S.; STARICHKOVA, V.Ye.

Effect of the vibration milling on the hydrolysis of polysaccharides of millet hulls. Izv.vys.ucheb.zav.;pishch.tekh.
no.5:105-109 '58. (MIRA 11:12)

1. Odesskiy tekhnologicheskii institut imeni I.V.Stalina,
kafedra organicheskoy khimii.
(Millet) (Polysaccarides) (Hydrolysis)

STARICHKOVA, V.Ye.

Effect of different grinding methods on the ~~hydrolysis of~~
the polysaccharides of millet husk. Izv. vys. ucheb. zav.;
~~piishch.tekh.~~ no.2:22-26 '60. (MIRA 14:7)

1. Odesskiy tekhnologicheskii institut imeni I.V. Stalina.
(Millet)
(Polysaccharides)
(Hydrolysis)

STARICHKOVA, V.Ye.; DUDKIN, M.S.; GLADNEVA, A.N.; MAKSIMENKO, N.S.

Preparation of fodder yeast from millet hulls. Gidroliz. i lesokhim.
prom. 16 no.1:9-11 '63. (MIRA 16:2)

1. Odesskiy tekhnologicheskii institut im. M.V.Lomonosova (for Starichkova, Dudkin).
 2. Krasnodarskiy gidroliznyi zavod (for Gladneva, Maksimenko).
- (Yeast as feeding stuff)

STARHA, Franci, ing. (Licki Osik)

Pin bearings today. Stroj vest 7 no. 4-5:100-104 0 61.

1. Tvornica MOL, Licki Osik..

STARHA, Franci, inz. (Licki Osik, AS-3)

Reconstruction of roller bearings for the lubricants of railroad cars. Tehnika Jug 17 no.11:Suppl.: Masinstvo 11 no.11:2109-2119 N '62.

1. Konstruktor kotrljajucih lezajeva u tvornici "Mol", Licki Osik.

STARIHA, Franci, dipl. inz. strojnistva

Selection of roller bearings. Stroj vest 9 no.4/5:109-116 0 '63.

1. BELT, Crnomelj.

STARIHA, Franci, inz., konstruktor kotrljajucih lezista (Licki
Osik AS-3, SR Hrvatska)

Analysis of the pressing process in the manufacture of rings
and races for antifriction bearings. Tehnika Jug 18 no.10:
Supplement: Masinstvo 12 no.10:1873-1881 0'63.

1. Tvornica "Mol", Licki Osik.

STARINA, Franci, inz.

Control in the production of ball bearings. Nova proizv 15
no.1/2:68-80 '64.

Starik, A.M.

57-9-22/40

AUTHORS: Golubchin, G.N., Starik, A.M.

TITLE: The Dependence of the Efficiency of the Auxiliary Discharge in Broad-Band Dischargers Upon the Position of the Ignition Electrode (Zavisimost' effektivnosti vspomogatel'nogo razryada v shirokopolosnykh razryadnikakh ot polozheniya elektroda podzhiga)

PERIODICAL: Zhurnal Tekhn. Fiz., 1957, Vol 27, Nr 9, pp. 2089-2091 (USSR)

ABSTRACT: The dependences of the ignition losses and energy losses at the peak (maximum) of broad-band dischargers of the ten centimeter range on the position of the ignition electrode are determined. The character of these dependences is explained by the non-uniform electron density distribution according to the length of the glowing discharge. The maximum of ignition losses and the energy minimum of the peak correspond to such an electrode position in which the high-frequency discharge space is filled by the glowing luminescence. An estimate of the distance between the cathode and the domain of glowing luminescence carried out at the conditions of this experiment was 0,4 to 0,5 mm. There are 4 figures and 1 Slavic reference.

SUBMITTED: December 4, 1956

AVAILABLE: Library of Congress
Card 1/1

SOV/109-3-11-6/13

Influence of the Higher-order Waves on the Characteristics of a System of Resonant Irises

in Figures 3 - 10. Figure 3a shows the frequency characteristic of a system in which diaphragms were spaced at a large distance; Figure 3b shows the characteristic for the case of $l/\lambda_0 = 0.128$. Figure 4 shows the value of the standing wave ratio as a function of l/λ_0 while Figure 5 illustrates the dependence of the interaction coefficient α on l/λ_0 ; from Figure 5, it is seen that for $l/\lambda_0 > 0.38$, the effect of the higher-order waves is negligible. Figures 6 and 7 illustrate the dependence of the standing wave ratio and α on λ_0/λ_{kp} , where λ_{kp} is the critical wavelength. Figure 8 shows the frequency characteristics of a system having $\lambda/\delta_0 = 0.32$ and $\lambda_0/\lambda_{kp} = 0.653$. The dependence of the standing wave ratio on the Q of the irises is illustrated in Figure 9, while the dependence of α on l/λ_0 for various values of Q is shown in Figure 10. From these experimental results,

Card2/3

SOV/109-3-11-6/13

Influence of the Higher-order Waves on the Characteristics of a
System of Resonant Irises

it is concluded that the higher-order wave interaction can be neglected provided the two irises are spaced at a distance of $l/\lambda_B > 0.25$, where λ_B is the length of the fundamental wave in the waveguide. The authors thank S.A. Sergeyev and S.I. Rudkovskiy for collaboration in carrying out the experiments. There are 10 figures and 2 references, 1 of which is Soviet and 1 English.

SUBMITTED: October 29, 1957

Card 3/3

81156

S/109/60/005/07.001/024
E140/E163

9.2140

AUTHOR: Starik, A.M.

TITLE: ~~Principal Directions~~ in the Development of Antenna
TR-Switches (Review)

PERIODICAL: Radiotekhnika i elektronika, Vol 5, No 7, 1960,
pp 1035-1051 (USSR)

ABSTRACT: The article presents a survey of American, English,
French and Russian work in this field between 1946 and
1958. Principal emphasis is given to integrated systems and
plug-in units (inserts) described in Ref 40. The majority of
Russian work cited concerns theoretical studies of bandwidth,
energy considerations, microwave gas-discharge, etc. See also
article on pp 1124-1128 of the present journal. HT
There are 15 figures and 69 references, of which 31 are English,
11 French and 27 Soviet.

SUBMITTED: May 6, 1959, and after revision, October 10, 1959.

Card 1/1

S/109/60/005/07/011/024

E140/E163

Application of Glow Discharge to Electronic Tuning of Transmitter
Blocking Discharger

obtained in the device. An electronic tuning band of $\pm 20\%$ may be obtained with easily obtainable values of electron concentration. The maximum loss in the 3-cm band should not exceed 1.5 dB. The necessary concentrations may be obtained using the cathode portion of a glow discharge. Using plasma, with an appreciably lower electron concentration, the length of the device must be increased substantially.

There are 2 figures, 2 tables and 7 references, of which 1 is Soviet and 6 are English.

SUBMITTED: November 21, 1959

Card 2/2

28528

S/109/61/006/009/011/018
D201/D302

9.3150 (also 1049, 1502, 1482)

AUTHOR: Starik, A.M.

TITLE: Propagation of microwaves in a waveguide containing
cathode parts of a glow discharge

PERIODICAL: Radiotekhnika i elektronika, v. 6, no. 9, 1961,
1433 - 1539

TEXT: The present paper presents certain data obtained from the study of model waveguides with cold cathodes of considerable length inside them. The experimental sections of the waveguide are shown diagrammatically in Fig. 1. The first had the cathode in the shape of a circular wire 0.4 mm diameter, stretched at an angle to the waveguide to provide correct matching. The second construction had the shape of a rhombic plate 0.4 mm thick. Both ends of the waveguide had very low q glass resonant windows. After proper evacuation, the final gas pressure was established and the phase shift and attenuation experimentally measured as functions of neon and

Card 1/5

28528

S/109/61/006/009/011/018
D201/D302

Propagation of microwaves in ...

helium pressure in the waveguide. The phase shift was measured from the shift of the mode of the standing wave. Results of various models were within 10 % of each other. Slightly larger than for the phase, the results of attenuation determination were due to incomplete evacuation of the device, showing more in the attenuation than in the phase shift. The results of measurements of attenuation L and phase shift θ as functions of neon pressure in plane cathode models for current $I = 12$ mA are shown graphically. The overall area of the cathode was 8.65 cm^2 with the resulting cathode current density $j_k = 1.4 \text{ mA/cm}^2$. The SWVR of the waveguides $10 \times 23 \text{ mm}$ did not exceed 1.15 in the absence of discharge. Similar in character the dependence of L and θ on neon pressure was found in waveguides $4 \times 23 \text{ mm}$. Comparative results for different waveguide dimensions are tabulated, giving the pressures corresponding to maxima of curves (p_{\max}) and the phase shift values of the maxima for every case. The interelectrode distance h used to be determined as the distance between the internal surface of the wide wall of the waveguide and the nearest surface to it, of the cathode $h =$

Card 2/5

28528

S/109/61/006/009/011/018

D201/D302

Propagation of microwaves in ...

$(b - t)/2$ where t - thickness of cathode. It is shown that P_{max} as determined from the attenuation curve, varies inversely proportional to the interelectrode distance. Figures show the dependence of the phase shift attenuation of discharge to potential U_p decrease on pressure at various points of discharge, for waveguides filled with neon and having a cylindrical cathode, the dependence on pressure of the quantity $F = 100(L/\theta)$, where L is in db and θ in degrees, F as function of pressure with helium-filled waveguides. It is stated in conclusion that the phase shift and attenuation produced by cathode parts of the glow discharge in rectangular waveguides exhibits sharp maxima with varying pressure. This maximum is in the region where $v_{cal} \approx \omega$ (v_{cal} - frequency of collisions).

These maxima could be explained by the shift of the virtual cathode surface, as the result of which the effective interaction between SHF oscillations and plasma is reduced for pressures greater than P_{max} . The smaller displacement of the maximum of the phase shift

Card 3/5

STARIK, A.M.

Cathode field of an anomalous glow discharge in a waveguide.
Radiotekh. i elektron. 10 no.4:779-780 Ap '65. (MIRA 18:5)

L 1170-66

ACCESSION NR: AP5017661

UR/0109/65/010/007/1250/1251
621.372.852.2/.3

40
B

AUTHOR: Starik, A. M.

TITLE: Estimating the density of electrons in a hollow-cathode gas-discharge attenuator

SOURCE: Radiotekhnika i elektronika, v. 10, no. 7, 1965, 1250-1251

TOPIC TAGS: attenuator, gas discharge attenuator

ABSTRACT: Operation of an original gas-discharge attenuator, in which the entire internal surface of the waveguide acts as a hollow cathode and an anode is mounted externally, is considered. An experimental attenuation-per-cm-length vs. cathode-current-density curve for a 7.2x3.4-mm waveguide is shown. An electron density of 1.6×10^{13} per cm^3 , at a He pressure of 50 torr, and a cathode-current density of 32 ma/ cm^2 is estimated. Orig. art. has: 2 figures and 5 formulas.

ASSOCIATION: none

SUBMITTED: 22May64

NO REF SOV: 004

Card 1/1 AP

ENCL: 00

OTHER: 004

SUB CODE: EC

USSR/Analytical Chemistry - Analysis of Inorganic Substances, G-2

Abst Journal: Referat Zhur - Khimiya, No 1, 1957, 123⁴

Author: Starik, I. Ye., Starik, A. S., Lozhkina, G. S., and Talitskaya, L. V.

Institution: Academy of Sciences USSR

Title: A Method for the Determination of Ionium

Original
Periodical: Byul. komis. po opredeleniyu absolyut. vozrasta geol. formatsiy AN
SSSR, 1955, Vol 1, 47-52

Abstract: After dissolution of the resin in HNO_3 the Th isotopes are deposited on Ce (carrier) as the oxalates. RaD, RaE, and Po are separated by electrolysis in 1 N HNO_3 by passing a 100 ma, 2.1 v current through the solution for 9 hours. UX_1 is used as an indicator for the completeness of Io separation. It has been established that: (1) Complete removal of Ra and U is achieved by double deposition of Ce(Io) oxalate; (2) the deposit of Ce oxalate after double deposition adsorbs 7-12% Po, >30% RaE, and 2-3% RaD; and (3) when H_2S is utilized to separate Ce(Io) from RaD, RaE, and RaF, complete separation is

Card 1/2 APPROVED FOR RELEASE: 08/25/2000

CIA-RDP86-00513R001652910020-5"

USSR/Analytical Chemistry - Analysis of Inorganic Substances, G-2

Abst Journal: Referat Zhur - Khimiya, No 1, 1957, 123⁴

Abstract: achieved, with the adsorption, however, of 30% of the I₂ on the sulfide precipitate.

STARIK, I.Ye.; RATNER, A.P. [deceased]; GROSHEV, G.V.; MURIN, A.N.;
STARIK, A.S.; GREBENSHCHIKOVA, V.I.; KLOKMAN, V.P.; NEPEDOV, V.D.;
LUR'YE, B.G.; ISHINA, V.A.; SMIRNOV, L.A.; YEFIMOVA, Ye.I.;
TOROPOVA, M.A.; SIMONYAK, Z.N.; FRENKLIKH, M.S.; SHCHEMELEVA, Ye.V.,
redaktor; VODOLAGINA, S.D., tekhnicheskiy redaktor

[A collection of practical studies in radio chemistry] Sbornik
prakticheskikh rabot po radiokhimi. [Leningrad] 1956. 210 p.
(MIRA 10:1)

1. Leningrad. Universitet.
(Radiochemistry)

STARIK, I.Ye.; STARIK, A.S.; YASHUGINA, Ye.A.; SMIRNOVA, Ye.A.

Quantitative separation of actinium from radioactinium and
actinium.X. Trudy Radiev.inst.AN SSSR. 8:170-176 '58.
(MIRA 12:2)

(Actinium--Analysis)

STARIK, A.S.; LEONT'YEV, V.G.

Method for determining microdoses of cesium from biological samples.
Vop. med. khim. 7 no.5:537-539 S-0 '61. (MIRA 14:10)

1. The I.M.Sechenov Institute of Evolutional Physiology of the
Academy of Sciences of the U.S.S.R.
(CESIUM-ANALYSIS)

SHAKHIDZHANYAN, L.G.; STARIK, A.S.; FLEYSIMAN, D.G.; GLAZUNOV, V.V.;
LEONT'YEV, V.G.; NESTEROV, V.P.

Distribution of radioactive cesium and strontium in human and
animal organs. Izv. AN SSSR. Ser. biol. no.3:442-448 My-Je '62.
(MIRA 15:6)

1. Institute of Evolutionary Physiology, Academy of Sciences
of the U.S.S.R., Leningrad.

(CESIUM--ISOTOPES)

(STRONTIUM--ISOTOPES)

(RADIOISOTOPES--PHYSIOLOGICAL EFFECT)

STARIK, D.E., kand.tekhn.nauk

Preliminary determination of the cost of engines.

Trudy MAI no. 161:47-59 '62.

(MIRA 15:12)

(Airplanes--Engines--Cost)

ANDRIANOV, D.P., doktor ekon. nauk, prof.; GENDEL'MAN, M.Z.,
kand. tekhn. nauk, dots.; GLICHEV, A.V., kand. ekon.
nauk, dots.; DIDENKO, S.I., kand. ekon. nauk, dots.;
ZHURAVLEV, A.N., kand. tekhn.nauk, prof.; ZAKHAROV,
K.D., kand. tekhn.nauk, dots.; MOISEYEV, S.V., kand.
tekhn. nauk, dots.; OL'SHEVETS, L.M., kand. tekhn.
nauk, dots.; ORLOV, N.A., prof.; POPOV, P.G., ispolnya-
yushchiy obyazannosti dots.; SARKISYAN, S.A., kand. ekon.
nauk, dots.; STARIK, D.E., kand. tekhn.nauk, ispolnyayu-
shchiy obyazannosti dots.; TER-MARKARYAN, A.N., kand.
tekhn. nauk, prof.; TIKHOMIROV, V.I., kand. tekhn.nauk,
prof.; CHESNOKOV, V.V., kand. ekon. nauk, dots.;
SHERMAN, Ye.I., kand. ekon. nauk, dots.; EL'BERT, L.M.,
kand. ekon. nauk, dots.; LAPSHIN, A.A., dots., retsenzent;
NOVATSKIY, V.F., kand. ekon. nauk, red.; TUBEYANSKAYA, F.G.,
red. izd-va; KARPOV, I.I., tekhn. red.

[Organization, planning and economics of airplane produc-
tion] Organizatsiia, planirovanie i ekonomika aviatsionnogo
proizvodstva. [By] D.P.Andrianov i dr. Moskva, Oborongiz,
1963. 694 p. (MIRA 16:10)

(Airplane industry--Management)

8/2535/63/000/154/0070/0080

ACCESSION NR: AT4031065

AUTHOR: Starik, D. E. (Candidate of Technical Sciences); Smirnova, A. P. (Engineer);
Yegorov, V. M. (Engineer)

TITLE: The planning of work according to the experimental-structural theme

SOURCE: Moscow. Aviatsonnyy institut. Trudy*, no. 154, 1963. Ekonomicheskaya
effektivnost' aviatsionnoy tekhniki (economic efficiency in aeronautical engineering),
70-80

TOPIC TAGS: economic efficiency, expenditure, experimental structural theme,
operation planning, calendar planning

ABSTRACT: The authors shed light on some questions of preliminary determination of
the expenditures on an experimental-structural theme and associated problems of
operation-calendar planning for experimental production. The basis for the initia-
tion of the experimental-structural work includes: the naming of products, the
object for which the product is intended, the product customer, the neighboring
organizations, the amount of products sent to the customer and the completion per-
iods, and sources of financing (state budget or self-support). The results were

Card 1/2

ACCESSION NR: AT4031065

presented in graphs and tables. The authors also determined the work capacity of fulfilling the stages and substages, and showed the composition of the planned calculation. Orig. art. has: 3 figures and 3 tables.

ASSOCIATION: Moscow Aviatseyonnyy institut (Moscow Institute of Aeronautics)

SUBMITTED: 00

DATE ACQ: 16Apr64

ENCL: 00

SUB CODE: AD

NO REF SOV: 000

OTHER: 000

Card 2/2

STARIK, Fye

CA

Radio content of petroleum waters of the Island Chikhen. N. V. TAGERVA AND P. F. STARIK. *Compt. rend. acad. sci. U. R. S. S. 1931A, 163 7.* -The waters containing petroleum in the Island Chikhen contain $4.37 \pm 0.75 \times 10^{-11}$ g. of Ra as sol. Ra salts per liter. For the detn. of Ra content a standardized universal electroscope was used, supplemented by ionization chamber. (P. F. STARIKOV)

ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION

RESEARCH

RESEARCH

STARIK, F.YE.

Relative leaching of uranium and radium isotopes from
uraninite. I. B. Starik, F. Y. Starik, and E. P. Koryakov.
Byull. Komissii Opredele. Absolyut. Vozrastu Geol. Formatsii,
Akad. Nauk S.S.S.R. 1953, No. 1, 22-23. — From 1 to 6-g.
samples of uraninite particles (dimension 0.5-1.0 mm.)
contg. 54.3 ± 0.3 wt. % U, $1.3 \times 10^{-4} \pm 0.06$ g. Ra/g.
mineral, and 2.3 ± 0.1 wt. % Th were leached with HNO_3 ,
 H_2O , and Na_2CO_3 at room temp. for one week. The amts.
of leached U and Ra in 0.1, 0.01, 0.001, 0.0001N HClO_4 ,
distd. water, and 0.001, 0.01, 0.1N Na_2CO_3 were, resp.:
 11 ± 1.5 , 11.2 ± 0.8 ; 1.6 ± 0.1 , 3.5 ± 0.05 ; $0.007 \pm$
 0.001 , 0.2 ± 0.015 ; 0.007 ± 0.001 , 0.1 ± 0.01 ; $0.004 \pm$
 0.0005 , 0.075 ± 0.005 ; 0.03 ± 0.007 , 0.016 ± 0.002 ; $0.4 \pm$
 0.07 , 0.09 ± 0.003 ; and 1.8 ± 0.1 , 0.25 ± 0.05 wt. %.
With respect to U, as data show, Ra was leached prefer-
entially in every case. A. P. Koryakov

nu Sci
60

3

PMF nyc

STARIK, F. Ye.

Fluoride method of separating small amounts of uranium
and its subsequent polarographic determination / I. E.
Starik, P. E. Starik, and A. N. Apollonova. *Trudy*
Khimicheskogo Inst. im. V. G. Khlopina; Khim. i Geokhim. 7,
107-10 (1954).—A new method is described which permits
one to sep. small quantities of U as the fluoride from other
elements: Th compds. act as a carrier. The fluorides of
Ca, rare earths, and Pb are copptd. with that of U; however,
these elements do not interfere with the subsequent polaro-
graphic detn. of U. A procedure of detg. polarographically
U in the presence of Th salts has been established. The
above method of analysis of small amts. of U is suitable for
its detn. in nature. Alfred Krembeller.

27 8
4E4
4E2C
4E3d
//

pm
mt

Starik F.E.

27
7
1-AE2C
1-4E4C
New aspect of analysis of small amounts of Cr^{+++} based on the determination of the extent of extinction of luminescence of uranium. I. E. Starik, F. E. Starik, and G. N. Kostyrev. *Trudy Radiovoego Inst. im. V.G. Khlopina, Khim. i Geokhim.* 7, 111-113 (1956); cf. following abstr. — Small amounts of Cr (down to 10^{-4} g.) can be detected by its quenching action of luminescence of U in a NaF head. At 8×10^{-4} g. U is present in the NaF head, the luminescence intensity is diminished by the addn. of more than 10^{-4} g. of Cr^{+++} or 10^{-3} g. of CrO_3 ; complete extinction of luminescence is observed with addns. above 10^{-3} g. A. K.

SR MT

STARIK, F. E.

27
Influence of various elements on the luminescence of uranium in sodium fluoride. I. E. Starik, L. E. Starik, L. Ya. Atrashenok, G. B. Kostyrev, V. N. Kozlov, and A. Ya. Krylov. *Trudy Radiofiz. Inst. im. V. G. Kalashina, Akim. Geokhim.* 7, 114-23 (1956); cf. *C.A.* 51, 17573e. The influence of 46 elements upon the luminescence of U in NaF has been studied. According to their behavior, the elements can be essentially arranged in 5 classes: (1) Na, K, Rb, Zn, Ti, S, Mo, W, Cl, Br, and I do not exhibit a marked influence, even if they are present in large aunts.; (2) Be, La, Th, P, Ni, Fe, Mn, Cu, Sr, Cd, Mg, B, Se, Cs, Zn, Ba, Li, and Si cause quenching of luminescence, if they are present in aunts. of several % of the wt. of the NaF bead; (3) Ag, Hg, Pb, Bi, Cr, and Co are strongly quenching elements, if their quantity is several tenths of 1% of the wt. of the bead; (4) Ca, Al, Ti, and Sn induce an increase in luminescence intensity or a change in its color; and (5) Ce, V, Nb, Ta, and Sb exhibit their own emission in NaF. Some elements show double action depending on their concentration; for instance, the luminescence is enhanced at a certain concentration, while it is quenched if the concentration is increased. It is suggested that the determination of uranium should be feasible in some liquids, without previous separation, by luminescence analysis. 29 literature references. Alfred Kerenheller

10
4E3d
4E4c

MT

STARIK, F. Ye.

STARIK, I. Ye.; STARIK, F. Ye.

Chromatographic analysis of small quantities of lead. Trudy Radiov.
inst. AN SSSR 5 no.2:129-133 '57. (MLRA 10:8)
(Lead) (Chromatographic analysis)

STARIK, F. YE.,

Starik, F. Ye., A. N. Yelizarova - Comparative Leaching Out of Several Isotopes.

The Sixth Session of the Committee for Determining the Absolute Age of Geologic Formations at the Department of Geologic-Geographical Sciences (OGGN) of the USSR Academy of Sciences at Sverdlovsk in May 1957

Izv. Ak Nauk SSSR, Ser. Geol., No. 1, 1958, p. 115-117 author Pekarskaya, T. B.

STARIK, I.Ye.; STARIK, F.Ye.; YELIZAROVA, A.N.; PLEKHADEV, Ye.P.

Leaching AcI from minerals. Biol.Kor. po opr.abs.vost.gool.forn.
no.3:60-61 '50. (MIRA 12:11)
(Leaching) (Radio--Isotopes)

STARIK, F. Ye.

AUTHORS: Starik. I. Ye., Starik. F. Ye., Apollonova, A. N. 78-1-23/43

TITLE: Adsorption of Micro Quantities of Uranium by Ferric Hydroxide and Desorption by Means of the Carbonate-Method. (Adsorbtsiya mikrokolichestv urana gidrookis'yu zheleza i desorbtsiya yego karbonatnym metodom).

PERIODICAL: Zhurnal Neorganicheskoy Khimii, 1958, Vol. 3, Nr 1, pp. 121-128 (USSR).

ABSTRACT: The adsorption of uranium on iron is important for analytical chemistry, since iron is often used as a carrier substance. The authors used U²³³ in their investigations. First the adsorption on ferric hydroxide with increasing pH is investigated. Carbonate-free ammonia serves here as basis. The maximum in the curve between pH 5 and pH 8 is explained by the fact that the hydroxide colloids are charged with the same signs outside of this range. This was electrophoretically proved. In carbonate solution the curve shows first a similar course which, however, declines steeply after pH 5,3, since uranium dissolves as complex carbonate and iron precipitates completely. The precipitation in ammoniacal medium was investigated with various quantities of uranium and iron with respect to its completeness. The precipitations and their results are summarized in a table.

Card 1/3

Adsorption of Micro Quantities of Uranium by Ferric Hydroxide
and Desorption by Means of the Carbonate Method.

78-123/43

Prior to their dealing with desorption, the authors investigate the influence of the alkali carbonates, especially of the ammonium carbonate, on the precipitation of the iron. Further the desorption of uranium is investigated, quantities of 10^{-5} g are quantitatively desorbed, with quantities of 10^{-6} - 10^{-8} g, however, losses up to 35% occur. This is attributed to the penetration of uranium into glass or platinum with the evaporation of the acid solution, as is proved. These losses can be avoided by adding metatitanic acid.

Conclusions:

- 1) The coprecipitation of micro quantities of uranium with ferric hydroxide takes place by adsorption.
- 2) The capability of adsorption depends on the pH of the solution in ammoniacal and carbonate solution.
- 3) Micro quantities of uranium precipitate with metatitanic acid under certain conditions.
- 4) The conditions of desorption of micro quantities of uranium (10^{-5} g - 10^{-8} g) from ferric-hydroxide-colloid (U:Fe = 1:10⁵) were determined by the carbonate-method.
- 5) The conditions of complete separation of micro quantities of uranium (10^{-5} - 10^{-8} g) from solutions by means of adsorption with ferric hydroxide were determined.

Card 2/3

Adsorption of Micro Quantities of Uranium by Ferric Hydroxide
and Desorption by Means of the Carbonate-Method.

78-1-23/43

There are 5 figures, 8 tables, and 24 references, 13 of which are Slavic.

SUBMITTED: June 18, 1957.

AVAILABLE: Library of Congress.

Card 3/3

SOV/7-58-5-7/15

AUTHORS: Starik, I. Ye., Starik, F. Ye.,
Mikhaylov, B. A.

TITLE: On the Problem of the Shift of Isotopic Ratios in Natural
Formations (K voprosu o smeshchenii izotopnykh sootnosheniy v
prirodnykh obrazovaniyakh)

PERIODICAL: Geokhimiya, 1958, Nr 5, pp. 462 - 464 (USSR)

ABSTRACT: The method suggested by V.V.Cherryntsev (Refs 8,9) makes use
of the measurement of the alpha and beta activity for the
determination of the U II - U I ratio. The small β -activity
can, however, be measured only with a low accuracy: therefore
the authors of this article modified this method. As U II
has a considerably shorter half life than U I the U II amount
may be neglected and the total amount of uranium may be taken
as measuring standard for U I. The sum of U I and U II is
determined by the alpha activity. Polonium was electrolytically
separated in the radiochemical purification, the thorium iso-
topes were separated by the precipitation with cerium fluoride
and radium isotopes by the precipitation with barium sulfate.
Aluminium and iron were separated by means of ammonium carbonate.
Uranium was determined by weighing. Uranium was separated

Card 1/3

On the Problem of the Shift of Isotopic Ratios
in Natural Formations

SOV/7-58-5-7/15

from a 0,4 m ammonium oxalate solution onto a target and the alpha activity was determined by means of an apparatus of the type AA. By means of this method some minerals of different age were investigated (Table 1): uraninite, pitchblende, uranium pitch ore, and schroekingerite (Shrekengerit). Only the last mentioned, which is a quaternary formation, showed a deviation of the isotopic ratio of uranium. Furthermore the effect of the leaching out with HNO_3 and Na_2CO_3 on the isotopic ratio was investigated (Table 2). An effect was demonstrated only in the leaching out of uraninite by means of Na_2CO_3 . At present the authors of this article investigate the kinetics of the sublimation of uranium from pitchblende. The isotopic ratio of sublimated uranium (800°) was determined. There are 2 tables and 11 references, 8 of which are Soviet.

ASSOCIATION: Radiyevyy institut im.V.G.Khlopina AN SSSR, Leningrad (Leningrad Radium Institute imeni V.G.Khlopina, AS USSR)

Card 2/3

On the Problem of the Shift of Isotopic Ratios
in Natural Formations

SOV/7-58-5-7/15

SUBMITTED: July 22, 1957

Card 3/3

STARIK, I.Ye.; NIKOLAYEV, D.S.; STARIK, F.Ye.; MELIKOVA, O.S.

Uranium content in natural waters of the U.S.S.R. Report No. 1.
Trudy Radiev.inst. AN SSSR. 8:250-261 '58. (MIRA 12:2)
! (Uranium) (Water--Analysis)

STARIK, I.Ye.; STARIK, F.Ye.; APOLLONOVA, A.N.

Carbonate method for separating microquantities of uranium from
iron. Trudy kom.anal.khim. 9:264-273 '58. (MIRA 11:11)
(Uranium) (Iron)

SOV/75-14-3-9/29

5(2)

AUTHORS:

Starik, I. Ye., Starik, F. Ye.,
Lazarev, K. F.

TITLE:

Photometric Determination of Micro-Quantities of
Thorium (Fotometricheskoye opredeleniye mikrokolichestv
toriya)

PERIODICAL:

Zhurnal analiticheskoy khimii, 1959, Vol 14, Nr 3,
pp 306-312 (USSR)

ABSTRACT:

The optical conditions for the photometric determination of thorium were devised on the basis of standard curves by means of the colorimetric photometer FEK-M using thoron as reagent. As can be seen from the figure the influence exercised by Ce and La upon the light absorption is suppressed at pH 0.96 - 0.85. Small calcium amounts do not disturb. The separation of thorium from sodium, potassium, calcium, and barium is carried out by precipitation of thorium together with $\text{Fe}(\text{OH})_3$. The quantitative precipitation was checked with UX_1 and RdTh . The separation of thorium from iron and uranium was carried out in weakly acid solution by precipitation with calcium oxalate. The mean absolute error was $\pm 0.3\%$ at $1 - 10 \mu\text{g Th}$.

Card 1/2

Photometric Determination of Micro-Quantities of
Thorium

SOV/75-14-3-9/29

The maximum error does not exceed $\pm 0.5\%$. There are 1 figure,
4 tables, and 23 references, 5 of which are Soviet.

ASSOCIATION: Radiyevyy institut AN SSSR, Leningrad imeni V. G. Khlopina
(Institute of Radium imeni V. G. Khlopin, Academy of Sciences,
USSR, Leningrad)

SUBMITTED: February 3, 1958

Card 2/2


S/186/61/003/006/008/010
EO40/E185

AUTHORS: Starik, I.Ye., Starik, F.Ye., and Yelizarova, A.N.

TITLE: Direct determination of protactinium and actinium
in uranites

PERIODICAL: Radiokhimiya, v.3, no.6, 1961, 749-754

TEXT: Detailed knowledge of the relative concentrations of individual radioisotopes in the various radioactive series of elements is absolutely essential in interpretation of radioactive dating data obtained especially by the lead technique. In case of the actinide series, the radioactive equilibrium between Pa^{231} , Ac^{227} and U^{235} can be determined by a direct measurement only, because indirect methods pre-suppose a priori that such an equilibrium already exists. As a continuation of the previously undertaken investigations of the authors on the radiochemistry of uranites (lead dating and separation of isotopes), a direct determination was made of protactinium and actinium in samples of the same mineral, using methods reported previously (Ref.9; I.Ye. Starik, A.P. Ratner, M.A. Pasvik, L.D. Sheydina, ZhAKh, Card 1/ 2



Direct determination of protactinium.. S/186/61/003/006/008/010
EO40/E185

v.12, 1, 87, 1957. Ref.10: I.Ye. Starik, L.D. Sheydina, ZnHKh, v.3, 1, 140, 1958). It was found that radioactive equilibrium exists between protactinium and actinium in well preserved specimens of uranites. Because of this, the authors find it difficult to generalise the results to include various weakly-radioactive minerals. D.M. Ziv and Ye.A. Volkova are mentioned in connection with their contributions in this field. There are 6 tables and 16 references; 10 Soviet-bloc and 6 non-Soviet-bloc. The English language references read as follows:
Ref.6: A.G. Maddock, G.L. Miles, J.Chem.Soc., s.i., v.2, 248, 1949.
Ref.7: A.V. Grosse, M.S. Agruss, J.Am.Chem.Soc., v.56, 10, 2200, 1934.
Ref.8: A. Grosse, J.Am.Chem.Soc., v.52, 5, 1742, 1930.

SUBMITTED: August 3, 1960

Card 2/2

S/081/62/000/004/014/087
B149/B101

AUTHORS: Starik, I. Ye., Starik, F. Ye., Yelizarova, A. N.
TITLE: Comparative leaching properties of some isotopes
PERIODICAL: Referativnyy zhurnal. Khimiya, no. 4, 1962, 113, abstract
4G15 (Byul. Komis. po opredeleniyu absolyutn. vozrasta geol.
formatsiy, AN SSSR, no. 14, 1961, 160-165)

TEXT: Investigation has been made of leaching the isotopes of Ra, Th, and Pb from specimens of uraninites (from Chkalov and Kamennaya Taybola mines) and monazite (Alakurti). The methods of determination used were as follows: Th by colorimetry; Ra, ThX, AcX, RdTh, RdAc, UX₁ and Ac - radiochemically; Pb - electrolytically; the isotope analysis of Pb by mass-spectrometry. The leaching of Ra isotopes (Ra²²⁶, ThX and AcX) has been carried out in 0.1 N HNO₃ from the demolished and intact specimens of uraninite. The demolished specimen showed larger percentage of leaching, and in both specimens AcX and ThX > Ra. The leaching from uraninites of Th has been carried out in 0.1 and 0.01 N HNO₃, 0.1 N

Card 1/2

STARIK, I.Ye.; STARIK, F.Ye.; YELIZAROVA, A.N.

Determination of protactinium and actinium in uraninite. Biul.Kom
po opr.abs.vozr.geol.form. no.5:72-75 '62. (MIRA 15:11)
(Uraninite) (Geological time)

3

PROCESSES AND PROPERTIES INDEX

Adorption of radium by glass. J. E. STARR and A. M. GURRICH. *Compt. rend. acad. sci. (U. R. S. S.)* 1931A, No. 12: 2871-5. Radioactivity of neutral and acid soln. in glass tubes contg. 3.6×10^{-4} g. of Ra in 5 cc. of soln. was measured with a Schmitt app. The initial activity of the neutral soln. was 211 divisions per min.; of the acid soln. (N/6 HCl), 297.2 div./min. After 20 days the radioactivity became practically const., having decreased in the acid soln. to 210 and in the neutral to 145. Addn. in the beginning to the neutral soln. of HCl to a concn. of N/6 raised the activity to 295.4. Addn. of HCl after the neutral soln. reached its min. radioactivity raised the activity to 210. Ra is thought to be absorbed and adsorbed by glass. Absorption is independent of acidity of soln. but the adsorbed layer is washed off by acid. FELIX P. LEE

METALLURGICAL LITERATURE CLASSIFICATION

ASB-SLA

GROUP 1

SECTION 1

SECTION 2

SECTION 3

SECTION 4

SECTION 5

SECTION 6

SECTION 7

SECTION 8

SECTION 9

SECTION 10

SECTION 11

SECTION 12

SECTION 13

SECTION 14

SECTION 15

SECTION 16

SECTION 17

SECTION 18

SECTION 19

SECTION 20

SECTION 21

SECTION 22

SECTION 23

SECTION 24

SECTION 25

SECTION 26

SECTION 27

SECTION 28

SECTION 29

SECTION 30

SECTION 31

SECTION 32

SECTION 33

SECTION 34

SECTION 35

SECTION 36

SECTION 37

SECTION 38

SECTION 39

SECTION 40

SECTION 41

SECTION 42

SECTION 43

SECTION 44

SECTION 45

SECTION 46

SECTION 47

SECTION 48

SECTION 49

SECTION 50

SECTION 51

SECTION 52

SECTION 53

SECTION 54

SECTION 55

SECTION 56

SECTION 57

SECTION 58

SECTION 59

SECTION 60

SECTION 61

SECTION 62

SECTION 63

SECTION 64

SECTION 65

SECTION 66

SECTION 67

SECTION 68

SECTION 69

SECTION 70

SECTION 71

SECTION 72

SECTION 73

SECTION 74

SECTION 75

SECTION 76

SECTION 77

SECTION 78

SECTION 79

SECTION 80

SECTION 81

SECTION 82

SECTION 83

SECTION 84

SECTION 85

SECTION 86

SECTION 87

SECTION 88

SECTION 89

SECTION 90

SECTION 91

SECTION 92

SECTION 93

SECTION 94

SECTION 95

SECTION 96

SECTION 97

SECTION 98

SECTION 99

SECTION 100

CA

Colloidal properties of polonium. I. E. Starik and L. V. Kozlov. *Trav. Inst. des radium* (U. S. S. R.) 2, 91-102 (in German 108-2)(1953); cf. C. A. 28, 910. --Parchment adsorbs Po better from an acid than from a basic medium. Min. adsorption is obtained from a neutral medium. Adsorption is greater when salts are present. Bleed charcoal adsorbs Po from a dil. acid soln. 100%. Best from NH₄OH soln. adsorption is min. All this is explained on the basis of colloidal distribution of Po in a basic or neutral medium. S. L. Madorsky

ASD-31A METALLURGICAL LITERATURE CLASSIFICATION

CA

Determination of radium in rocks and minerals by the emanation method. L. E. Starik and A. S. Sinagina. *Trav. inst. fiz. radium* (U.S.S.R.) 2, 104 (1954 English transl. 1955).—Methods were worked out for dissolving rock and mineral samples preliminary to measuring their emanation intensities. The general plan consisted in dissolving the sample then adsorbing the Ra from solution on BaCl₂ by first pptg. the Ba as a sulfate and then converting the sulfate to a chloride by dissolving in HCl. The methods were much simplified to save time. In all cases iron or porcelain dishes were used instead of Pt. The Ra contents were detd. in torbernite, khalopinite, zircon and grandfleurite and of mixts. of khalopinite, monazite and feldspar and of zircon, monazite and diopside. S. L. Madorsky

CA

PROCESSES AND PROPERTIES INDEX

A study of possible errors in α -particle measurements. J. E. Starik and O. S. Melikova. *Trav. inst. fiz. radium* (U.S.S.R.) 2, 117-20 (in English 120) (1933).—The cause of variation in α -ray measurements was studied. Most of the variations and errors are due to the type of vessel used for holding the sample. All metal vessels give concordant results but glass dishes can be used under some conditions. Other nonconducting vessels, such as porcelain or paraffin show variations reaching as high as 800%. S. L. M.

ASME-SLA METALLURGICAL LITERATURE CLASSIFICATION

3

ca

A criticism of the use of the photographic method for the investigation of the colloidal state of polonium. I. E. Starik and M. Yu. Deizenrot-Muisovskaya. *Compt. rend. acad. sci. U. R. S. S. (N. S.)*, 1, 540-3 (in German 543-6) (1934).—Exptl. evidence is given to prove that the photographic method is unsuitable for the investigation of the colloidal state of radioactive elements because of the adsorption of these elements on mica, vaseline, etc., used to coat the photographic plate. M. McMahon

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND ORDERS

PROCESSES AND PROPERTIES INDEX

8

CA

The geochemistry of uranium and radium. I. E. Starik. *Abad. V. I. Vernadskomu Pyatidesyatiletiyu Nauch. Dopysheni* 1, 445-62(1936); *Chem. Zentr.* 1936, I, 1565. The leaching out of U and Ra from minerals by ground water is discussed. Expts. show that Ra is more readily extd. by neutral, weakly acid or weakly alk. media than U. It is assumed that Ra is not built into the crystal lattice of the U-contg. minerals but is present rather as an inclusion in the capillaries of the mineral. U, on the other hand, is built into the crystal lattice. In this way the difference in the geochem. behavior of the 2 elements is explained. The migration of U is therefore greatly retarded since it can occur only with soln. of the U-contg. mineral. It can be expected, however, that U once in soln. is more readily held in soln. than other elements, especially Ra, which can be transported only short distances in soln. In secondary minerals the radioactive equil. is destroyed because of the migration of Ra.

M. G. Moore

ASB-55A METALLURGICAL LITERATURE CLASSIFICATION

1A

1ST AND 2ND ORDERS

PROCESSES AND PROPERTIES INDEX

2ND AND 4TH ORDERS

3

Radiochemical analysis of khlopinite. I. Composition and constitution of khlopinite. I. E. Starik and N. M. (1947) *Dokl. Akad. Nauk SSSR* 13, 211-12 (in English 217) (1947). From the relations obtained for the groups of oxides ($M_2O_3 \cdot MO_2 \cdot MO = 1:1:2$) present in a mineral from Khilok (Transbaikalia) there is assumed the existence of a new $Cb-Ti$ mineral. The mineral, analyzed as $M_2O_3 \cdot Cb_2O_3 \cdot TiO_2$, has been named "khlopinite" in honor of V. G. Khlopin. II. Leaching of radium and uranium from khlopinite. *Ibid.* 218-26 (in English 226-7). -- Expts. on the leaching of Ra and U from khlopinite reveal that the leaching of Ra in neutral, slightly acid, and alk. media is considerably greater than that of U. John Livak

ASB-3LA METALLURGICAL LITERATURE CLASSIFICATION

1947-1948

1949-1950

1951-1952

1953-1954

1955-1956

1957-1958

1959-1960

1961-1962

1963-1964

1965-1966

1967-1968

1969-1970

1971-1972

1973-1974

1975-1976

1977-1978

1979-1980

1981-1982

1983-1984

1985-1986

1987-1988

1989-1990

1991-1992

1993-1994

1995-1996

1997-1998

1999-2000

2001-2002

2003-2004

2005-2006

2007-2008

2009-2010

2011-2012

2013-2014

2015-2016

2017-2018

2019-2020

2021-2022

2023-2024

2025-2026

2027-2028

2029-2030

2031-2032

2033-2034

2035-2036

2037-2038

2039-2040

2041-2042

2043-2044

2045-2046

2047-2048

2049-2050

2051-2052

2053-2054

2055-2056

2057-2058

2059-2060

2061-2062

2063-2064

2065-2066

2067-2068

2069-2070

2071-2072

2073-2074

2075-2076

2077-2078

2079-2080

2081-2082

2083-2084

2085-2086

2087-2088

2089-2090

2091-2092

2093-2094

2095-2096

2097-2098

2099-2100

2101-2102

2103-2104

2105-2106

2107-2108

2109-2110

2111-2112

2113-2114

2115-2116

2117-2118

2119-2120

2121-2122

2123-2124

2125-2126

2127-2128

2129-2130

2131-2132

2133-2134

2135-2136

2137-2138

2139-2140

2141-2142

2143-2144

2145-2146

2147-2148

2149-2150

2151-2152

2153-2154

2155-2156

2157-2158

2159-2160

2161-2162

2163-2164

2165-2166

2167-2168

2169-2170

2171-2172

2173-2174

2175-2176

2177-2178

2179-2180

2181-2182

2183-2184

2185-2186

2187-2188

2189-2190

2191-2192

2193-2194

2195-2196

2197-2198

2199-2200

2201-2202

2203-2204

2205-2206

2207-2208

2209-2210

2211-2212

2213-2214

2215-2216

2217-2218

2219-2220

2221-2222

2223-2224

2225-2226

2227-2228

2229-2230

2231-2232

2233-2234

2235-2236

2237-2238

2239-2240

2241-2242

2243-2244

2245-2246

2247-2248

2249-2250

2251-2252

2253-2254

2255-2256

2257-2258

2259-2260

2261-2262

2263-2264

2265-2266

2267-2268

2269-2270

2271-2272

2273-2274

2275-2276

2277-2278

2279-2280

2281-2282

2283-2284

2285-2286

2287-2288

2289-2290

2291-2292

2293-2294

2295-2296

2297-2298

2299-2300

2301-2302

2303-2304

2305-2306

2307-2308

2309-2310

2311-2312

2313-2314

2315-2316

2317-2318

2319-2320

2321-2322

2323-2324

2325-2326

2327-2328

2329-2330

2331-2332

2333-2334

2335-2336

2337-2338

2339-2340

2341-2342

2343-2344

2345-2346

2347-2348

2349-2350

2351-2352

2353-2354

2355-2356

2357-2358

2359-2360

2361-2362

2363-2364

2365-2366

2367-2368

2369-2370

2371-2372

2373-2374

2375-2376

2377-2378

2379-2380

2381-2382

2383-2384

2385-2386

2387-2388

2389-2390

2391-2392

2393-2394

2395-2396

2397-2398

2399-2400

2401-2402

2403-2404

2405-2406

2407-2408

2409-2410

2411-2412

2413-2414

2415-2416

2417-2418

2419-2420

2421-2422

2423-2424

2425-2426

2427-2428

2429-2430

2431-2432

2433-2434

2435-2436

2437-2438

2439-2440

2441-2442

2443-2444

2445-2446

2447-2448

2449-2450

2451-2452

2453-2454

2455-2456

2457-2458

2459-2460

2461-2462

2463-2464

2465-2466

2467-2468

2469-2470

2471-2472

2473-2474

2475-2476

2477-2478

2479-2480

Determination of the age of rocks by the radioactive method. I. New method for the determination of small amounts of lead and its use to determine the age of khlopinite. I. E. Sharik and N. M. Segel. *Trav. inst. elat radium* (U.S.S.R.) 3, 280 (in English 210) (1937).— A new method, based on copptn. with isomorphous salts, has been developed for the sepn. of Pb in khlopinite. with slight modification it can be applied to most minerals. The most favorable conditions for the copptn. of Pb with BaSO_4 , are studied. Pb is best sepd. from BaSO_4 by electrolysis in uranyl nitrate soln. The colorimetric method for detn. of Pb is not sufficiently accurate to det. the age of rocks and minerals. The age of khlopinite as detd. by the Pb method is 175×10^6 yr., and 127×10^6 yr. by the He method. Such good agreement indicates that ordinary Pb is probably absent and that the mineral had not undergone any appreciable changes. J. L.

1ST AND 2ND ORDERS

PROCESSES AND PROPERTIES INDEX

CA

Adsorption of radium by glass. I. E. Starik and A. M. Gurevich. *Trav. inst. chim. radium* (U.S.S.R.) 3, 241-56 (in English 256) (1937); cf. C. A. 26, 4535. —The extent of Ra adsorption by glass from RaBr₂ soln. contg. 5×10^{-3} g. Ra per cc. apparently depends on the amt. of impurities present. In a carefully purified soln. the degree of adsorption remains const. in the range pH 6.6-4.6. Further increase in acidity sharply lowers the adsorption, which becomes practically negligible at pH 2.3. The data for slightly alk. media (pH 7.9) cannot be compared with those for acid media because of the possible presence of colloid particles (SiO₂, etc.) in the alkali used. Change in adsorption with time is greatly dependent upon the impurities present in the soln. and is probably due to the presence of complex colloid particles on which Ra is adsorbed.

John Livak

COMMON ELEMENTS

COMMON VARIETIES INDEX

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

SECTION 1

SECTION 2

SECTION 3

SECTION 4

SECTION 5

SECTION 6

SECTION 7

SECTION 8

SECTION 9

SECTION 10

SECTION 11

SECTION 12

SECTION 13

SECTION 14

SECTION 15

SECTION 16

SECTION 17

SECTION 18

SECTION 19

SECTION 20

SECTION 21

SECTION 22

SECTION 23

SECTION 24

SECTION 25

SECTION 26

SECTION 27

SECTION 28

SECTION 29

SECTION 30

SECTION 31

SECTION 32

SECTION 33

SECTION 34

SECTION 35

SECTION 36

SECTION 37

SECTION 38

SECTION 39

SECTION 40

SECTION 41

SECTION 42

SECTION 43

SECTION 44

SECTION 45

SECTION 46

SECTION 47

SECTION 48

SECTION 49

SECTION 50

SECTION 51

SECTION 52

SECTION 53

SECTION 54

SECTION 55

SECTION 56

SECTION 57

SECTION 58

SECTION 59

SECTION 60

SECTION 61

SECTION 62

SECTION 63

SECTION 64

SECTION 65

SECTION 66

SECTION 67

SECTION 68

SECTION 69

SECTION 70

SECTION 71

SECTION 72

SECTION 73

SECTION 74

SECTION 75

SECTION 76

SECTION 77

SECTION 78

SECTION 79

SECTION 80

SECTION 81

SECTION 82

SECTION 83

SECTION 84

SECTION 85

SECTION 86

SECTION 87

SECTION 88

SECTION 89

SECTION 90

SECTION 91

SECTION 92

SECTION 93

SECTION 94

SECTION 95

SECTION 96

SECTION 97

SECTION 98

SECTION 99

SECTION 100

19

8

The inconstancy of the lead ratio in different parts of the mineral chlopinite. I. E. Starik and O. S. Melikova. *Trav. inst. Etal radium* (U. S. S. R.) 4, 384-42 (in English, 1952-53) (1958). --In various fractions of a homogeneous chlopinite which has been broken up, the Ra and Th vary by 25% but the Pb varies as much as 55%. Thus different ages are calcd. for the fractions. Probably more l and Th than Pb have been leached from the mineral H. M. Leicester

ASB SLA METALLURGICAL LITERATURE CLASSIFICATION

Dr. A. J. L.

Migration of ionium under natural conditions. I. B. Starik and O. S. Belikova
(Compt. rend. Acad. Sci. U. R. S. S., 1941, 31, 911-913).--The I° content of carno-
tite with an isomorphous admixture of tyuyamunite has been determined. Although the
migration of I° from the mineral is considerable, the I° content is relatively near
its equilibrium val.
A. J. L.

DR. Chemical Sci. Geochemistry. Mbr., Radiochemical Lab., All-Union Inst.,
Leningrad, -1941-.

STARIK, I. Ye.

"Comparative Migration Capacity of Radium and Mesothorium: I. A Finding of Ferrithorit in North Kirghizia," Dokl. AN, 32, No 4, 1941.

11-13, 1960, 1961, 1962

Dr. A. A.

Age of phroxenite intrusions of Afrikanda and Ozernaja Varaca in the Kola Peninsula. A. K. Gerling and I. E. Starik (Compt. rend. Acad. Sci., U. R. S. S., 1962, 25, 153-154).--Two shorlowite specimens from the above intrusions were investigated, their content of He, Ra, and Th being determined. The two intrusions were formed simultaneously $\sim 340 \times 10^6$ years ago. A. J. C.

PRINCIPLES AND PROPERTIES INDEX

3

4

Radiological study of Caucasian region mineral waters
I. E. Stark. *Bull. acad. sci. U.R.S.S., Class. sci. chim.*
1943, 435-42 (English summary).—The enrichment of
mineral waters by Ra and Rn was studied; previously,
the following factors were not estd. at their true value:
contact time of water with the rocks, filtration speed of
water, water-capacity and porosity of the stratum and the
state of preservation of the stratum. G. M. Kozlovskii.

ASAC SEA DETAIL/DECHAL LITERATURE CLASSIFICATION

Form of approach and the conditions for the primary migration of radioelements in nature. I. E. Statik. *Izvestiya Khim.* 12, 287-307(1963).—Review with 46 references, largely to Russian literature. Copptn. and absorption are considered as factors leading to the enrichment of elements found only in very low concns. 20 tables and figures. F. H. Rathmann

ASA SLA BIBLIOGRAPHICAL LITERATURE CLASSIFICATION

STARIK, I. Ye., Order Labor Red Banner, 1945.

"On the Geochemistry of the Carboniferous and Permian Deposits of the
Chistopol Region of Tataria," Dokl AN, 49, No 9, 1945. Cor. Mbr., Acad. Sci.,
-1947-, Consultant, Wismath AG, Germany.

ISSN/Acad Sci
Chem - Acad Sci

Aug 1947

Radium

"June Session of Department of Chemical Sciences"
1 1/2 pp

"Vest Akad Nauk SSSR" No 8

Session called in honor of 25th anniversary of the
Radium Institute of the Academy of Sciences. I. Ye.
Starik, Acting Director of the Radium Institute,
read a paper on the work of the Institute. Twenty-
eight workers awarded prizes and medals: Academician
V. G. Khlopkin, Director of the Institute; Academician
P. I. Lukirevskiy; B. A. Nikitin, I. Ye. Starik,
5779

USSR/Acad Sci (Contd)

Aug 1947

and A. A. Grinberg, Corresponding Members of the
Academy of Sciences; M. G. Meshcheryakov, M. A.
Pavlik-Khlopina, and A. Kh. Ratner, Candidates in
Chemical Sciences, etc. Several scientists, among
them A. N. Nesmeyanov and A. F. Kapustin, sub-
mitted articles and papers for judgment.

Mbr., Sci. Council, -1947-
Deputy Dir., -1947-

STARIK, I. YE.

5779

CP

* Vitalii Grigor'evich Khlopov. B. A. Nikitin and I. E.
Starik. *Izvest. Akad. Nauk S.S.S.R., Otdel. Khim. Nauk*
1950, 121-6.—Biography, with portrait and review of
scientific work on 60th jubilee. G. M. K.

1. STARIK, I.Ye.
2. USSR (600)
4. Geological Time
7. Current state of radioactive methods of determining the age of old and young formations, Izv.AN SSSR. Ser.geol. no. 4, 1952.

9. Monthly List of Russian Accessions, Library of Congress, APRIL 1953, Uncl.

STARIK, I. YE.

PA 241T43

USSR/Geophysics - Radioactive Methods Nov/Dec 52

"Contemporary State of Radioactive Methods for the
Determination of Age of Old and Young Formations,"
I. Ye. Starik

"Iz Ak Nauk SSSR, Ser Geol" No 6, pp 11-20

Presents committee's report, given at a session of
Dept of Geologico-Geographical Sci, Acad Sci USSR,
on 12-13 Apr 52, on detn of the abs age of geological
formations. Concludes that present methods are
unsatisfactory and require more persistent, collec-
tive efforts by geologists, geophysicists, physicists,
and chemists.

241T43

(CA 47 no. 14: 6837 13)

1. STARIK, I. YE.; RATNER, A. P.
2. USSR 600
4. Radioactivity
7. "Chemistry of radioactive elements," collected works. Reviewed by
I. YE. Starik, A. P. Ratner, Sov. kniga, No. 1, 1953.

9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

1. IIA IK, I. Ye.
2. USSR (600)
4. Radioactivity
7. Current state of radioactive methods of determining the age of old and young formations. Izv. AN SSSR. Ser. geol. No. 6, 1953.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

STARIK, I.Ye., otvetstvennyy redaktor; SHCHERBAKOV, D.I., akademik, redaktor; VINOGRADOV, A.P., akademik, redaktor; BARANOV, B.I., professor, redaktor; GERLING, E.K., professor, redaktor; LEVIN, B.Yu., kandidat fiziko-matematicheskikh nauk, redaktor; KRYLOV, A.Ya., redaktor; PEKARSKAYA, T.B., kandidat geologo-mineralogicheskikh nauk; MYASNIKOV, I.A., redaktor; POLYAKOVA, T.V., tekhnicheskii redaktor.

[Transactions of the first session of the Commission on Determining the Absolute Age of Geologic Formations] Trudy pervoi sessii komissii po opredeleniiu absolutnogo vozrasta geologicheskikh formatsii; 12-15 aprelia 1952 g. Moskva, Izd-vo Akademii nauk SSSR, 1954. 231 p. (MLR 8:1)

1 Chlen-korrespondent Akademii nauk SSSR (for Starik). 2. Akademiya nauk SSSR. Otdeleniye geologo-geograficheskikh nauk.
(Earth--Age)

SP. F. L. T.
IOFFE, A.F.; LEBEDEV, A.A.; FOK, V.A.; STARIK, I.Ye.; KONSTANTINOV, B.P.;
DZHELEPOV, B.S.; PERFILOV, N.A.; DOBRETISOV, L.N.; STARODUBTSEV, A.V;
NEMILOV, Yu.A.; ZHDANOV, A.P.; MURIN, A.N.; AGLINTSEV, K.K.; TSARE-
VA, T.V.; SHUL'MAN, A.R.; YEREMEYEV, M.A.

P.I.Lukirskii; obituary. Vest.AN SSSR 24 no.12:62 D '54.(MIRA 8:1)
(Lukirskii, Petr Ivanovich, 1894-1954)

STARIK. I.Ye., redaktor; SHCHERBAKOV, D.I., akademik, redaktor; VINOGRADOV, A.P., akademik, redaktor; POLKANOV, A.A., akademik, redaktor; SHATSKIY, N.S., akademik, redaktor; BARANOV, V.I., professor, redaktor; PEKARSKAYA, T.B., kandidat geologo-mineralogicheskikh nauk, redaktor; CHERDYNTSEV, V.V., redaktor; POLYAKOVA, T.V., tekhnicheskij redaktor.

[Transactions of the third session of the Committee for Determining the Absolute Age of Geological Formations, March 25-27, 1954] Trudy tret'ei sessii, 25-27 marta 1954. g. Moskva, 1955. 260 p. [Microfilm] (MLRA 9:1)

1. Akademiya nauk SSSR. Komissiya po opredelniya absolyutnogo vozrasta geologicheskikh formatsii. 2. Chlen-korrespondent AN SSSR (for Starik). (Geological time)

STARIK, I.Ye

Relative leaching of uranium and radium isotopes from
uraninite. I. B. Starik, P. S. Starik, and E. P. Kozlov.
Byull. Komissii Uranium. Akad. Nauk S.S.R. 1955, No. 1, 29-32. — From 1 to 6 g.
samples of uraninite particles (dimension 0.5-1.0 mm.)
contg. 54.3 ± 0.3 wt. % U, $1.6 \times 10^{-7} \pm 0.06$ g. Ra/g.
mineral, and 2.3 ± 0.1 wt. % Th were leached with HNO_3 ,
 H_2O , and Na_2CO_3 at room temp. for one week. The amts.
of leached U and Ra in 0.1, 0.01, 0.001, 0.0001N HNO_3 ,
distd. water, and 0.001, 0.01, 0.1N Na_2CO_3 were, resp.:
 11 ± 1.5 , 11.2 ± 0.8 ; 1.6 ± 0.1 , 3.5 ± 0.05 ; $0.007 \pm$
 0.001 , 0.2 ± 0.016 ; 0.007 ± 0.001 , 0.1 ± 0.01 ; $0.004 \pm$
 0.0005 , 0.075 ± 0.006 ; 0.03 ± 0.007 , 0.016 ± 0.002 ; $0.4 \pm$
 0.07 , 0.09 ± 0.003 ; and 1.2 ± 0.1 , 0.25 ± 0.05 wt. %.
With respect to U, as data show, Ra was leached prefer-
entially in every case. A. P. Kozlov

nu Sci
6/1

3

RML nje

Radium Inst. in V. G. Khlopun, A S USSR

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 1, 15-57-1-497
p 80 (USSR)

AUTHORS: Starik, I. Ye., Melikova, O. S., Kurbatov, V. V.,
Aleksandruk, V. M.

TITLE: The Relation of Temperature to the Emanation Factor of
Uraninite for Radon, Thoron, and Actinon (Zavisimost'
emaniruyushchey sposobnosti uraninita po radonu, toronu
i aktinonu ot temperatury)

PERIODICAL: Byul. Komis. po opredeleniyu absolyut. vozrasta geol.
formatsii AN SSSR, 1955, Vol 1, pp 33-38.

ABSTRACT: Uraninite containing $2.16 \cdot 10^{-7}$ g/g Ra, $2.39 \cdot 10^{-1}$ g/g
Th, and $8.0 \cdot 10^{-11}$ g/g Ac was studied for emanations of
radon, actinon, and thoron during heating. It was
found that the emanation factor for radon (K_{Rn}), equal
to 2.1, remains almost constant up to 90° . It decreases
sharply to 1.0 in the interval from 90° to 100° . The
emanation factor for thoron (K_{Th}) is 1.14 at 21° , and
it gradually decreases to 0.75 on heating to 100° . The

Card 1/2

STARIK, I.YE.

19
Distribution of radioelements in various parts of uraninite.
I. B. Starik, O. S. Melikova, and E. V. Sobolevich. Byull.
Akad. Nauk S.S.S.R. 1955, No. 1, 89-91. Analysis of
specimens taken from the outer surface, the mid-portion, and
the central cores of specimens of uraninite, for content
of U, Pb, Ra, Th X, and Ac X, resulted in the following
findings. In good, intact specimens the content of the 5
elements was const. through the thickness of the specimen,
while in cracked and deformed specimens Ra, Ac, and U de-
clined from the center to the periphery, while Th and Pb rose,
indicating a leaching process which removed Ra at higher rate
than U, Th, or Pb. Generally, peripheral parts of all speci-
mens tended to be higher in Th and Pb than central parts
and in the central portions the Ac/U ratio was 1.7×10^{-10}
instead of the normally expected 2.08×10^{-10} . Ra tends
to migrate to the environment more than does U or the
products of decompn. of actinouranium. The emanation
coeffs. of powders of all the specimens from all parts of a
specimen are of the same order of magnitude since diffusion

becomes less important. Isotopic compn. of Pb from cen-
tral, well-preserved parts of specimens indicates good agree-
ment with the local geol. indications of age; only Pb²¹⁰/Th
ratio leads to some disagreements. Isotope distribution in
the outer portions of the specimens invariably gave a higher
indication of age than indicated by other methods, although
Pb²¹⁰/Pb²⁰⁶ ratio tended to give low results of age estn.
C. M. Kosolapov

STARIK, I YE
USSR/Analytical Chemistry - Analysis of Inorganic Substances, G-2

Abst Journal: Referat Zhur - Khimiya, No 1, 1957, 1234

Author: Starik, I. Ye., Starik, A. S., Lozhkina, G. S., and Talitskaya, L. V.

Institution: Academy of Sciences USSR

Title: A Method for the Determination of Ionium

Original
Periodical: Byul. komis. po opredeleniyu absolyut. vozrasta geol. formatsiy AN
SSSR, 1955, Vol 1, 47-52

Abstract: After dissolution of the resin in HNO_3 the Th isotopes are deposited on Ce (carrier) as the oxalates. RaD, RaE, and Po are separated by electrolysis in 1 N HNO_3 by passing a 100 ma, 2.1 v current through the solution for 9 hours. UX_1 is used as an indicator for the completeness of Io separation. It has been established that: (1) Complete removal of Ra and U is achieved by double deposition of Ce(Io) oxalate; (2) the deposit of Ce oxalate after double deposition adsorbs 7-12% Po, >30% RaE, and 2-3% RaD; and (3) when H_2S is utilized to separate Ce(Io) from RaD, RaE, and RaF, complete separation is

Card 1/2

STARIK, I. Ye.

USSR/ Geology - Rock formation

Card 1/1 Pub. 46 - 6/21

Authors : Starik, I. Ye.; Murina, G. A.; and Krylov, A. Ya.

Title : Criteria of the suitability of minerals for determination of their age
 by the helium method

Periodical : Izv. AN SSSR. Ser. geol. 20/2, 67 - 71, Mar-Apr 1955

Abstract : By making use of the fact, established through research, that there is
 a definite connection between the form in which radio-active elements
 are present in a mineral and the preservation of helium in the mineral
 during geological times, the age of rocks was successfully determined.
 Twenty types of granite were subjected to this helium method to deter-
 mine their age. Four references: 1 USA and 3 Soviet (1933-1946).
 Tables; graphs.

Institution :

Submitted : December 1, 1954

STARIK, I. Ye.
USSR/Chemistry - Technical books

Card 1/1 Pub. 147 - 22/22

Authors : Starik, I. Ye.; Murin, A. N.; and Nefedov, V. D.

Title : Critique and bibliography

Periodical : Zhur. fiz. khim. 29/11, 2110-2110-2111, Nov 1955

Abstract : Critical review is presented of the book by An. N. Nesmeyanov, A. V. Lapitskiy and N. P. Rudenko, entitled, "Derivation of Radioactive Isotopes," published by Goskhimizdat (State Publication of Chemical Literature) in 1954.

Institution :

Submitted :

STARIK, I.Ye.; RATNER, A.P. [deceased]; GROSHKOV, G.V.; MURIN, A.N.;
STARIK, A.S.; GREBENSCHIKOVA, V.I.; KLOKMAN, V.P.; NEFEDOV, V.D.;
LUR'YE, B.G.; ISHINA, V.A.; SMIRNOV, L.A.; YEFIMOVA, Ye.I.;
TOROPOVA, M.A.; SIMONYAK, Z.N.; FRENKLIKH, M.S.; SHCHEMENIEVA, Ye.V.,
redaktor; VODOLAGINA, S.D., tekhnicheskiy redaktor

[A collection of practical studies in radio chemistry] Sbornik
prakticheskikh rabot po radiokhimi. [Leningrad] 1956. 210 p.
(MIRA 10:1)

1. Leningrad. Universitet.
(Radiochemistry)

NIKITIN, Boris Aleksandrovich; STARIK, I.Ye., otvetstvennyy redaktor;
OKERBLUM, M.H., redaktor izdatel'stva; SMIRNOVA, A.V., tekhnicheskii redaktor

[Selected works] Izbrannye trudy. Moskva, Izd-vo Akademii nauk
SSSR, 1956. 349 p. (MIRA 9:11)

1. Chlen-korrespondent AN SSSR (for Nikitin, Starik)
(Chemistry, Physical and theoretical)
(Radioactivity)

FAUL, Henry, red.; AVDZEYKO, G.V.[translator]; STARIK, I.Ye.,
red.

[Nuclear geology] IAdernaia geologiia. Pod red. i s predisl.
I.E.Starika. Moskva, Izd-vo inostr.lit-ry, 1956. 555 p.
(MIRA 15:7)
(Geophysics) (Nuclear physics) (Radioactivity)

STARIK, I. YE.

USSR / Isotopes.

B-7

Abs Jour : Ref Zhur - Khimiya, No 8, 1957, 26052 .

Author : I.Ye.Starik, N.I. Alekseyenko, N.G. Rozovskaya

Title : Colloid Properties of Polonium

Orig Pub. : Izv Ak Nauk SSSR, Otdel Khim. Nauk, 745-63, 1957

Abstract : The possibility of existence of true colloids in very diluted solutions (10^{-13} to 10^{-9} M) was studied on the example of Po. The results of the study of the dependence on the solution concentration and pH of the processes of electrochemical separation of Po in Cu, adsorption and desorption on glass of various composition, centrifuging, ultrafiltration and radiography confirm the concepts of different states of Po in different media, proposed by the authors. At pH equal to 1 - 4, Po is in the ion non-hydrolyzed state; hydrolysis and the formation of positively charged colloidal particles starts at pH equal to 6 - 7; at pH equal to about 8 the recharge of particles takes place; insoluble Po compounds are formed and their equilibrium with

Card : 1/2

... compared from the results of the study of electrochemical separation of Po. It was established that the formation of colloids of radioactive elements did not depend on their specific properties connected with their radioactive nature.

Card : 2/2

STARIK, I. YE.

USSR / Isotopes.

B-7

Abs Jour : Ref Zhur - Khimiya, No 8, 1957, 26053

Author : I. Ye. Starik, N. I. Alekseyenko
Title : Radiographic Method of Studying State of Radioactive Elements in Diluted Solutions.

Orig Pub : Zh. teorg. khimii, 1, No 7, 1676 - 1679, 1956

Abstract : With a view to deciding upon the question of the applicability of the radiographic method to the study of the state of radioactive elements in diluted solutions, the character of the absorption of Po by photoemulsion, as well as its adsorption on mica and glass depending on the concentration of Po in the solution (1×10^{-11} to 5×10^{-8} M) and on pH (1.3 to 9.5) were investigated. Nuclear photographic plates of the A-2 type with an emulsion layer 60 μ thick were used. It was shown that the blackening of the photographic plate increased with the concentration rise of Po.

Card : 1/2

STARIK, I. Ye.

Role of secondary processes in the determination of rock ages by radioactive methods. I. B. Starik (Radium Inst., Acad. Sci. U.S.S.R., Leningrad). *Goskhatrazh* 1956, No. 5, 18-20. —The age of rocks was detd. by the Pb method. Minerals were studied for the emission and leachability of their different radioelements. Results of the emission studies showed that: (1) the coeff. of emissivity varies within wide limits, from 0.07 to 100%; (2) value of the coeff. of emissivity depends on the nature of the mineral and the degree of retentivity of a given sample; (3) value of the emissivity coeff. is a criterion of retentivity of a given sample; (4) on grinding the minerals, the emissivity changes very little for the most part; and (5) value of emission in water and in air usually differs negligibly (within limits of 10-15%). A study of the leachability of Ra, U, ThX, and RdTh showed: (1) radioelements, parents of the natural series, usually are leached less than are their decompn. products; (2) leachability of the decompn. products depends to a great extent on the cations present in soln., whereas cations present do not affect leachability of the parent substances; (3) per cent of leachability in neutral medium is usually less than the coeff. of emissivity; (4) cations may be arranged in a series according to the strength of their action on leachability of Ra isotopes: $H^+ > Ba^{++} > Pb^{++} > Sr^{++} > Ca^{++} > K^+ > Na^+$; and (5) effect of anions on leachability is related chiefly to acceleration or retardation of adsorption phenomena. Graphs comparing the leachability of Ra and U, and ThX and RdTh are presented. Results are given of age detn. of uraninites and monazite of Northern Karelia, radiochem. data for uraninites from pegmatites of Northern Karelia, emissivity coeffs. of some uraninite samples, radiochem. analyses of some samples of uraninite and surrounding rock, isotopic compn. of Pb from crust and central portion of a uraninite sample, per cent of radioelements leached from uraninite, and leachability of ThX, RdTh, Th, and U from monazite. 15 references.

3
1-RMB
1-gum

EE
RMB
RT

STARIK, I. Ye

Fluoride method of separating small amounts of uranium
and its subsequent polarographic determination. I. B.
Starik, B. B. Starik, and A. N. Apollonova. *Trudy
Khimicheskogo Inst. im. V. G. Khlopina, Khim. i Geokhim.* 7,
107-10 (1956).—A new method is described which permits
one to sep. small quantities of U as the fluoride from other
elements; Th compds. act as a carrier. The fluorides of
Ca, rare earths, and Pb are copptd. with that of U; however,
these elements do not interfere with the subsequent polaro-
graphic detn. of U. A procedure of detg. polarographically
U in the presence of Th salts has been established. The
above method of analysis of small amts. of U is suitable for
its detn. in nature. Alfred Krembeller

27 8
4E4
4E2C
4E3d
//

PM
mt

Starik, I. E.

27 7
New aspect of analysis of small amounts of Cr^{+++} based on the determination of the extent of extinction of luminescence of uranium. I. E. Starik, E. E. Starik, and G. B. Kostyrev. *Trudy Radiofiz. Inst. im. V. G. Khlopina, Khim. i Geokhim.* 7, 111-113 (1958); cf. following abstr. Small amounts of Cr (down to 10^{-4} g.) can be detected by its quenching action of luminescence of U in a NaF bead. If 8×10^{-4} g. U is present in the NaF bead, the luminescence intensity is diminished by the addn. of more than 10^{-4} g. of Cr^{+++} or 10^{-4} g. of CrO_4^{--} ; complete extinction of luminescence is observed with addns. above 10^{-3} g. A. K.

1-4E2C
1-4E4C

SR MT

15-1957-10-14145
Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 10,
p 124 (USSR)

AUTHORS: Starik, I. Ye., Starik, F. Ye., Atrashenkov, L. Ya.,
Kostyrev, G. B., Kosyakov, V. N., Krylov, A. Ya.

TITLE: The Influence of Different Elements on the Fluorescence
of Uranium in Sodium Fluoride (Vliyaniye razlichnykh
elementov na lyuminesentsiyu urana vo ftoristom natrii)

PERIODICAL: Tr. Radiyev. in-ta AN SSSR, 1956, vol 7, pp 114-125

ABSTRACT: The principal merits of the fluorescent method of deter-
mining U are its simplicity, speed, and high sensitivity.
It is possible to determine up to 1×10^{-10} grams of U
in the bead. The precision of the determination is gen-
erally close to 20%. In any method using different ac-
tivators, measuring devices, and sources of ultraviolet
light, impurities exert considerable influence by alter-
ing the fluorescence of the Uranium or by extinguishing
it. The influence of 45 elements on the fluorescence of
U in NaF was studied. The investigations were made on

Card 1/3

15-1957-10-14145

The Influence of Different Elements on the Fluorescence of Uranium in Sodium Fluoride

beads of NaF weighing 5 mg and containing 5×10^{-9} g of U. The elements were introduced into the bead either by mixing with NaF or with the corresponding salt, in different proportions, in case of soluble compounds, by dipping the bead of NaF, which contains a definite quantity of U, into the quenching salt solution. When using the dipping method, it is necessary to know the precise weight of the bead, inasmuch as beads of different weights take up different amounts of solution. The intensity of fluorescence is strongly influenced by the surface of the bead, which is a function of the quality of fusion, of the cooling of the bead, and also of the quantity of NaF. The elements investigated may be divided approximately into five groups, according to their influence on the fluorescence of uranium. 1) Na, K, Rb, Zn, Ti, S, Mo, W, Cl, Br, and J show essentially no effect, even in relatively large quantities. 2) Be, Ce, La, Th, P, Ni, Fe, Mn, Cu, Sr, Cd, Mg, B, Se, Cs, Zn, Ba, Li, and Si extinguish fluorescence when present in the bead in large quantities (on the order of several per cent of the weight of the bead). 3)

Card 2/3

15-1957-10-14145

The Influence of Different Elements on the Fluorescence of Uranium in Sodium Fluoride

Ag, Hg, Pb, Bi, Cr, and Co sharply extinguish the fluorescence when present in quantities several times ten per cent of the weight of the bead. 4) Ca, Al, Tl, and Sn strengthen the fluorescence or produce changes in the color. When the relative concentration of U to Ca is 1:10,000, U fluoresces green. With higher Ca content, the intensity of fluorescence begins to increase and then decrease. The fluorescent color changes from yellow-green to blue. When the concentration of Al is $5 \cdot 10^{-5}$ g in the bead, it does not fuse to transparency. Tl and Sn strengthen the fluorescence when their concentration in the bead is $1 \cdot 10^{-5}$ – $5 \cdot 10^{-5}$ g. 5) Ce, V, Nb, Ta, and Sb produce distinctive fluorescence in NaF. Ce and V give a red color, Sb a medium blue. The fluorescence of Nb and Ta is very similar to that for uranium. Some elements have a two-fold effect, depending on their concentration: with low concentrations they increase the fluorescence; with high they extinguish it.

Card 3/3

A. A. Rozbianskaya

STARIK, IYE.

19
Sampling of granites for radiochemical investigation.
I. E. Starik and A. Ya. Krylov. *Trudy Radiovogo Inst.*
im. V. G. Khlopina, Khim. i Geokhim. 7, 149-54 (1958).—
Suggestions and examples are given for fitting the type of
sampling to the size and character of the body being in-
vestigated and the knowledge desired. D. J. Milton
Geochemistry of radioactive elements in rocks of the
Kirovograd-Zhitomir magmatic complex of Ukraine. L.
V. Kozlov, M. S. Filippov, S. I. Danilevich, and K. S.
Ivanova. *Trudy Radiovogo Inst. im. V. G. Khlopina, Khim.*
i Geokhim. 7, 155-99 (1958).—Averages of widely varying
contents of 107 samples of these Archean granites are:
U 5.7×10^{-4} , Th 3.3×10^{-4} , and Ra $1.8 \times 10^{-4}\%$. Dif-
ferent intrusives vary markedly, but there is a correlation
with rock type. U is enriched in the border zones. High
Th content is often connected with a rare earth (monazite)
mineralization. D. J. Milton

8
1-1000
-1000

11
Rm

STARIK, Iye

27
Colloidal Properties of Polonium. I. E. Starik, N. I. Aleksenko, and N. G. Bozovskaya (Izv. Akad. Nauk S.S.S.R., 1958, (Khim.), (7), 755-763).—[In Russian].
 The question of colloidal behaviour of radioactive elements in infinitely dil. soln., and whether such elements themselves form true colloids or are adsorbed by other colloids always present in soln., is one of the least clarified problems of radiochemistry. A preliminary study was made of Po, the methods used being adsorption on and desorption from glasses, electrodeposition on Cu, centrifuging, ultrafiltration, all these applied to varying concentrations of Po, and autoradiography. The tabulated results of the tests, despite the fact that solubility valencies are approx., give a clear picture of their order and clarify the question of colloid-formation by the element itself. Since there is nothing sp. in the process, the methods adopted should prove equally efficient in the case of other elements (e.g. U, Ru, Ra, &c.). 17 ref.—Z. N. P.

6
 Met 1 PMZ
 1-JNM
 2 4E2C
 4E4j

fra pmt any
 xlll

STARIK, I. Ye.

27
 /Luminescent determination of uranium in rocks and minerals. I. B. Starik, L. Ya. Astrashenok, and A. Ya. Kevlov (V. G. Khlopin Radium Inst., Acad. Sci. U.S.S.R., Leningrad). *Geokhimiya* 1956, No. 8, 39-40. --The effect of different individual elements on the luminescence of U in NaF was studied. Also a study of several combinations of elements was made. The authors summarize the results obtained from these studies and report a further development of the method as applied to the study of different rocks and minerals. The method for detn. of U in rocks is as follows. The rock sample is fused with NaF and then removed from the crucible and pulverized. From the powder 5-mg. beads are prepd. for measurement of luminescence by means of a photometer. Accuracy of the measurements is $\pm 20\%$. Data obtained are presented in a table. Other tables show data from the detn. of U in minerals. With an adequate U content in rocks and minerals (usually 1×10^{-4} g./g.) the quant. luminescent detn. of U is applicable without sepn. from its accessory elements. For some rocks, rock-forming minerals, and most widely distributed accessory minerals, the concns. at which the effect of the elements was eliminated were detd. A comparison of results obtained by the method used and other methods gave entirely satisfactory results.

Gladys S. Macv

M fra
 any

STARIK, I. Ye.

Pierre Curie; on the fiftieth anniversary of his death. Izv. AN
SSSR. Ser. geol. 21 no. 4: 120-121 Ap '56. (MLRA 9:8)
(Curie, Pierre, 1859-1906)

STARIK, I YE.

USSR/Physical Chemistry - General Problems on Isotope Chemistry B-7

Abs Jour : Referat Zhur - Khimiya, No 2, 1957, 3676

Author : Starik I.Ye., Rozovskaya N.G.

Inst : Academy of Sciences USSR

Title : Study of State of Micro-Amounts of Radioelements by Desorption Method.

Orig Pub : Dokl. AN SSSR, 1956, 107, No 6, 850-852

Abstract : Investigation of desorption of Polonium (I), sorbed by different samples of glass from aqueous solutions of 0.1 N HNO₃, neutral, and 0.1 N NaOH. As desorbents were used 0.1 and 1 N HNO₃, distilled water and 0.1 N NaOH. Noted is the difference in behavior of I, sorbed from neutral and nitric acid containing media. The authors correlate this difference with the difference in the state of I in these media and consider that the method of desorption permits to form an opinion concerning the bonding strength of sorbed substance and surface and

Card 1/2

- 54 -

STAR-1K, 1V5

3
TE4
0

V. Determination of lead isotope composition in rocks.
E. Starik and E. M. Sobatovich. *Doklady Akad. Nauk S.S.S.R.* 111, 395-7(1958).—Pb was sepd. from rocks by volatilization in a furnace at 1400°. The vapors were reduced with H₂ and Pb was deposited on a cold surface. Preliminary tests showed that 80-90% of the Pb was recovered at a sublimation temp. of approx. 7/8 the m.p. of the rock. Pb was sepd. from the other volatile components by electrolysis of the nitrates. Pb was deposited on the anode. The anodal Pb(NO₃)₂ soln. was pptd. with KI and tested mass-spectrographically. Volatilization of Pb did not fractionate it isotopically. The accuracy of the method was confirmed by duplicate analyses of 2 granite samples. The isotope proportions found were Pb-206 = 17.5-19.4%, Pb-207 = 15.5%.

for
003

STARIKH, I. *YR*

*not in
file*
3776

7
DETERMINATION OF URANIUM IN THE ACCESSORY
MINERALS. I. E. Starikh, L. Ya. Atrachenok, and A. Ya.
Krylov (Khlopin Radium Inst.). Doklady Akad. Nauk S.S.S.R.
111, 644-6 (1956) Nov. 21. (In Russian) *3*

A method of direct fusion of U with a bead of NaF reveals
the quantities of U even in the tiniest crystals of accessory
minerals, without any preliminary chemical treatment.
Tables showing the estimations of minimum quantities of U
(in $\times 10^{-3}$ %) in mixture with other minerals. (R.V.J.)
DM m